

### **NHS GREATER GLASGOW AND CLYDE**

NB: This document should be read in conjunction with the current Summary of Product Characteristics (SPC)

## **DRUG AND INDICATION:**

Generic drug name:	Lamivudine	
Formulation:	Film-coated tablet containing 100mg lamivudine	
Intended indication:	Chronic hepatitis B infection in adults with compensated liver disease with	
	evidence of active viral replication, liver inflammation and/or fibrosis.	
Status of medicine or	Licensed medicine	
treatment:	Formulary medicine	

## RESPONSIBILITIES OF ACUTE CARE/SPECIALIST SERVICE (CONSULTANT):

- Undertake baseline investigations/monitoring and initiate treatment or ask GP to initiate treatment.
- If appropriate, ensure that the patient has an adequate supply of medication (usual minimum of 28 days) until the shared care arrangement are in place
- Dose adjustments

#### Acute care/specialist service will provide the GP with:

- An initiation letter (which includes diagnosis, relevant clinical information, treatment plan, duration of treatment before consultant review)
- Letter of outpatient consultations, ideally within 14 days of seeing the patient

### Acute care/specialist will provide the patient with relevant drug information to enable:

- Understanding of potential side effects
- Understanding of the role of monitoring

## RESPONSIBILITIES OF PRIMARY CARE (GENERAL PRACTITIONER):

- To prescribe in collaboration with the acute specialist according to this agreement
- To ensure the continuous prescription of medication until treatment is discontinued at specialist instruction
- Liaison with the hospital specialist in the event of symptoms or abnormal results thought due to this treatment

## RESPONSIBILITIES OF PATIENT:

- To attend hospital and GP clinic appointments. Failure to attend appointments may result in medication being stopped
- To report adverse effects to their specialist
- To request repeat prescriptions from the GP prior to current prescription finishing

## **ADDITIONAL RESPONSIBILITIES:**

None

SEPTEMBER 2019 SEPTEMBER 2022



### **NHS GREATER GLASGOW AND CLYDE**

### **CAUTIONS:**

- Renal impairment: dose adjustment recommended for patients if creatinine clearance < 50 ml/min, (see SPC).</li>
- Exacerbations of hepatitis
- Patients with decompensated liver disease
- Lactic acidosis
- Pregnancy and breastfeeding
- HIV co-infection. Used at higher dose with other anti-retroviral drugs

## **CONTRAINDICATIONS:**

Hypersensitivity to the active substance or to any of the excipients

### TYPICAL DOSAGE REGIMEN:

Route of administration:	Oral administration		
Recommended starting dose:	100mg every 24 hours with or without food		
Titration of dose:	None		
Maximum dose:	100 mg daily		
Conditions requiring dose adjustment:	Renal Impairment		
Usual response time:	Variable, depends on HBV viral load and host factors		
Duration of treatment	Treatment is usually for many years. Treatment may be		
	discontinued if there is HBsAg loss or HBeAg seroconversion.		

All dose adjustments or discontinuations will be decided in acute care and directions given in medical letter to the GP

## SIGNIFICANT DRUG INTERACTIONS:

Do not use with emtricitabine for HIV

# UNDESIRABLE EFFECTS:

ADR details	Management of ADR
Weakness, fatigue, headache,	These are the most common side-effects and usually mild
dizziness, nausea, vomiting, diarrhoea,	
abdominal pain.	
Muscle disorders, including elevations	
of CPK, myalgia and cramps	

The above list should not be considered exhaustive. For further documented ADRs and details of likelihood etc, see Summary of Product Characteristics or BNF.

## BASELINE INVESTIGATIONS (ACUTE SECTOR):

Urea and electrolytes, eGFR, LFTs, HIV test.

DOCUMENT PRODUCED BY: DOCUMENT APPROVED BY: DATE APPROVED: PLANNED REVIEW DATE: DR DAVID BELL, CONSULTANT IN INFECTIOUS DISEASES PRESCRIBING INTERFACE SUBCOMMITTEE OF ADTC SEPTEMBER 2019

SEPTEMBER 2019 SEPTEMBER 2022



### **NHS GREATER GLASGOW AND CLYDE**

## MONITORING (PRIMARY CARE):

No monitoring is to be undertaken in Primary Care

# MONITORING (ACUTE SECTOR):

The following monitoring is to be undertaken in Acute Care

Monitoring Parameters	Frequency	Laboratory results	Action to be taken
U&Es, LFTs	Every 3-6 months		
Hepatitis B Viral load	Every 3-6 months		
Hepatitis B e markers	Every 6 months		

### PHARMACEUTICAL ASPECTS:

- Shelf life 2 years
- Store below 30°C (tablets)

## Cost:

Approximate cost for 1 patient per year is £964 (BNF accessed online 22/8/19)

# INFORMATION FOR COMMUNITY PHARMACIST:

None of note

# **ACUTE CARE/SPECIALIST SERVICE CONTACT INFORMATION:**

Name	Designation	Acute Site	Department phone number
Dr David Bell	Consultant in Infectious	Brownlee Centre,	0141 301 7489
Dr Erica Peters	Diseases	Gartnavel General Hospital	
Dr Helen Cairns	Consultant	Cartnavel Conoral Hespital	0141 301 7489
Dr Matt Priest	Gastroenterologist	Gartnavel General Hospital	
Dr Stephen Barclay	Consultant	Classow Boyal Infirmary	0141 211 4911
Dr Ewan Forrest	Gastroenterologist	Glasgow Royal Infirmary	
		Queen Elizabeth University	0141 201 2177
Dr Judith Morris	Consultant	Hospital	
Dr Shouren Datta	Gastroenterologist		
		Victoria Infirmary	0141 347 8320
Dr Mathis Heydtmann	Consultant	Inverclyde Royal Hospital	01475 633 777
	Gastroenterologist		
	dustroenterologist	Royal Alexandra Hospital	0141 314 6850
Dr Rizwana Hamid	Consultant	Vale of Leven Hospital	01389 817 239
	Gastroenterologist	vale of Levell Hospital	
Kathryn Brown	BBV Specialist Pharmacists	Gartnavel General Hospital	0141 211 3383
Fiona Marra			0141 211 3317
Alison Boyle			

### **SUPPORTING DOCUMENTATION:**

DOCUMENT PRODUCED BY: DOCUMENT APPROVED BY: DATE APPROVED: PLANNED REVIEW DATE: DR DAVID BELL, CONSULTANT IN INFECTIOUS DISEASES PRESCRIBING INTERFACE SUBCOMMITTEE OF ADTC SEPTEMBER 2019

SEPTEMBER 2022

PAGE 3 OF 4



# **NHS GREATER GLASGOW AND CLYDE**

NHS GGC Hepatitis B Treatment Guideline
 http://www.staffnet.ggc.scot.nhs.uk/Info%20Centre/PoliciesProcedures/GGCClinicalGuidelines/GGC%20Clinic
 al%20Guidelines%20Electronic%20Resource%20Direct/Hepatitis%20B%20Infection%20Assessment%20and%2
 OManagement%20in%20Adult%20Patients.pdf

SEPTEMBER 2022